


**Order:** SAMPLE REPORT**Client #:** 12345**Doctor:** Sample Doctor, MD
Doctors Data Inc.
3755 Illinois Ave.
St. Charles, IL 60174**Patient:** Sample Report**Age:** 48**Sex:** Male**Body Mass Index (BMI):** N/A**Sample Collection** **Date/Time****Date Collected** 10/01/2018**AM30** 10/02/2018 0800**Night** 10/02/2018 2100**Date Received** 10/03/2018**Date Reported** 10/05/2018

Analyte	Result	Unit	L	WRI	H	Optimal Range	Reference Interval
Cortisol AM30	12	nmol/L		◆		14.0 - 25.0	5.1 - 30.0
Cortisol Night	1.2	nmol/L		◆		1.0 - 4.0	0.33 - 7.0
DHEA*	138	pg/mL		◆			137 - 336

Cortisol Graph**Hormone Comments:**

- The suboptimal AM cortisol level and reported symptoms are suggestive of HPA axis (adrenal gland) dysfunction.

Notes:

RI= Reference Interval, L (blue)= Low (below RI), WRI (green)= Within RI (optimal), WRI (yellow)= Within RI (not optimal), H (red)= High (above RI)

The current samples are routinely held three weeks from receipt for additional testing.

*This test was developed and its performance characteristics determined by Doctor's Data, Inc. The FDA has not approved or cleared this test; however, FDA clearance or approval is not currently required for clinical use. The results are not intended to be used as the sole means for clinical diagnosis or patient management decisions.

Methodology: Enzyme Immunoassay



Order: SAMPLE REPORT



Client #: 12345

Doctor: Sample Doctor, MD
Doctors Data Inc.
3755 Illinois Ave.
St. Charles, IL 60174

Patient: Sample Report

Age: 48

Sex: Male

Body Mass Index (BMI): N/A

Sample Collection Date/Time

Date Collected 10/01/2018

AM30 10/02/2018 0800

Night 10/02/2018 2100

Date Received 10/03/2018

Date Reported 10/05/2018

Analyte	Result	Unit	L	WRI	H	Reference Interval	Supplementation Range**
Estradiol (E2)	<0.5	pg/mL		◆		< 2.5	
Progesterone (Pg)	43	pg/mL		◆		< 94	500 - 3000
Pg/E2 Ratio	430				▲	200 - 300	
Testosterone	98	pg/mL		◆		30 - 143	110 - 500
DHEA*	138	pg/mL		◆		137 - 336	



Hormone Comments:

- The low Pg/E2 ratio is consistent with progesterone insufficiency (estrogen dominance), which may increase the risk of prostate gland enlargement and cancer. Supplementation with topical progesterone to correct this relative deficiency is a consideration.
- Suboptimal testosterone is consistent with reported deficiency symptoms and may be associated with metabolic syndrome (insulin resistance). Serum vitamin D, hemoglobin A1c and insulin levels may be warranted. Boosting the testosterone level is a consideration.

Notes:

RI= Reference Interval, L (blue)= Low (below RI), WRI (green)= Within RI (optimal), WRI (yellow)= Within RI (not optimal), H (red)= High (above RI)
The current samples are routinely held three weeks from receipt for additional testing.

The Pg/E2 ratio is an optimal range established based on clinical observation. Progesterone supplementation is generally required to achieve this level in men and postmenopausal women.

*This test was developed and its performance characteristics determined by Doctor's Data, Inc. The FDA has not approved or cleared this test; however, FDA clearance or approval is not currently required for clinical use. The results are not intended to be used as the sole means for clinical diagnosis or patient management decisions.

**If supplementation is reported then the supplementation ranges will be graphed. The supplementation ranges depicted are for informational purposes only and were derived from a cohort of adult men and women utilizing physiologic transdermal bioidentical hormone therapy.

Methodology: Enzyme Immunoassay