



Order: 999999-9999



Client #: 999999

Doctor: Sample Doctor, MD

Doctors Data Inc

123 Main St.

St. Charles, IL 60174 USA

Patient: Sample Patient

Id: 999999

Age: 38 DOB: 00/00/1986

Sex: Female

Sample Collection

Date Collected

Date/Time

05/14/2025

Date Received

05/23/2025

Date Reported

06/02/2025


Specimens Collected

3


Microbiome Abundance and Diversity Summary

The abundance and diversity of gastrointestinal bacteria provide an indication of gastrointestinal health, and gut microbial imbalances can contribute to dysbiosis and other chronic disease states. The GI360™ Microbiome Profile is a gut microbiota DNA analysis tool that identifies and characterizes more than 45 targeted analytes across six Phyla using PCR and compares the patient results to a characterized normobiotic reference population. The web chart illustrates the degree to which an individual's microbiome profile deviates from normobiosis.

LEGEND



The web image shows the relative diversity and balance among bacteria belonging to the six primary Phyla. The white shaded area represents the patient's results compared to a normobiotic reference population. The center of the web represents less abundance while the outer edges represent more than normobiotic.




Dysbiosis and Diversity Index

These indexes are calculated from the results of the Microbiome Profile, with scores ranging from 1 to 5, and do not include consideration of dysbiotic and pathogenic bacteria, yeast, parasites and viruses that may be reported in subsequent sections of the GI360™ test.

The Dysbiosis Index (DI) is calculated strictly from the results of the Microbiome Profile, with scores from 1 to 5. A DI score above 2 indicates dysbiosis; a microbiota profile that differs from the defined normobiotic reference population. The higher the DI above 2, the more the sample deviates from the normobiotic profile. The dysbiosis test and DI does not include consideration of dysbiotic and pathogenic bacteria, yeast, parasites and viruses that may be reported in subsequent sections of the GI360™ test.

A diversity score of 3 indicates an expected amount of diversity, with 4 & 5 indicating an increased distribution of bacteria based on the number of different species and their abundance in the sample, calculated based on Shannon's diversity index. Scores of 1 or 2 indicate less diversity than the defined normobiotic reference population.



GI Health Markers

Butyrate producing bacteria	<input type="checkbox"/>
Gut barrier protective bacteria	<input type="checkbox"/>
Gut intestinal health marker	<input type="checkbox"/>
Pro-inflammatory bacteria	<input checked="" type="checkbox"/>
Gut barrier protective bacteria vs. opportunistic bacteria	<input type="checkbox"/>

☒ = Expected ☐ = Imbalanced

Key Findings

Lactoferrin, High	% Valerate, Very Low	<i>Providencia alcalifaciens</i> , Cultured
Lysozyme, Very High	Butyrate, Low	<i>Candida krusei</i> , Cultured
Calprotectin, High	β-glucuronidase, Low	
% Acetate, Very High	<i>Citrobacter farmeri</i> / <i>amalonaticus</i> , Cultured	
% Propionate, Very Low	<i>Klebsiella oxytoca</i> , Cultured	
% Butyrate, Low	<i>Klebsiella pneumoniae</i> , Cultured	



Microbiome Bacterial Abundance; Multiplex PCR



Order: 999999-9999



Client #: 999999

Doctor: Sample Doctor, MD
Doctors Data Inc
123 Main St.
St. Charles, IL 60174 USA

Patient: Sample Patient

Id: 999999

Age: 38 **DOB:** 00/00/1986

Sex: Female

Sample Collection

Date Collected

Date/Time

05/14/2025

Date Received

05/23/2025


Date Reported

06/02/2025

Specimens Collected

3

LEGEND



Results are graphed as deviations from a normobiotic population. Normobiosis or a normobiotic state characterizes a composition of the microbiota profile in which microorganisms with potential health benefits predominate in abundance and diversity over potentially harmful ones.



Notes:

The gray-shaded area of the bar graph represents reference values outside the reporting limits for this test.

*This test was developed and its performance characteristics determined by Doctor's Data Laboratories in a manner consistent with CLIA requirements. The U. S. Food and Drug Administration (FDA) has not approved or cleared this test; however, FDA clearance is not currently required for clinical use. The results are not intended to be used as a sole means for clinical diagnosis or patient management decisions.

Methodology: Multiplex PCR

**Order:** 999999-9999**Client #:** 999999**Doctor:** Sample Doctor, MD
Doctors Data Inc
123 Main St.
St. Charles, IL 60174 USA**Patient:** Sample Patient**Id:** 999999**Age:** 38 **DOB:** 00/00/1986**Sex:** Female**Sample Collection****Date Collected****Date/Time**

05/14/2025

Date Received

05/23/2025

Date Reported

06/02/2025

Specimens Collected

3

**Notes:**

The gray-shaded area of the bar graph represents reference values outside the reporting limits for this test.

*This test was developed and its performance characteristics determined by Doctor's Data Laboratories in a manner consistent with CLIA requirements. The U. S. Food and Drug Administration (FDA) has not approved or cleared this test; however, FDA clearance is not currently required for clinical use. The results are not intended to be used as a sole means for clinical diagnosis or patient management decisions.

Methodology: Multiplex PCR



Microbiome Bacterial Abundance; Multiplex PCR



Order: 999999-9999



Client #: 999999

Doctor: Sample Doctor, MD
Doctors Data Inc
123 Main St.
St. Charles, IL 60174 USA

Patient: Sample Patient

Id: 999999

Age: 38 **DOB:** 00/00/1986

Sex: Female

Sample Collection

Date Collected 05/14/2025

Date Received 05/23/2025

Date Reported 06/02/2025


Specimens Collected 3

Firmicutes	Result	-3	-2	-1	0	+1	+2	+3	Reference Interval
<i>Streptococcus salivarius</i> ssp. <i>thermophilus</i>	0				▲				0
<i>Streptococcus</i> spp.	0				▲				0
<i>Veillonella</i> spp.	0				▲				0
Proteobacteria	Result	-3	-2	-1	0	+1	+2	+3	Reference Interval
Proteobacteria	+3							▲	0
<i>Enterobacteriaceae</i>	+3							▲	0
<i>Escherichia</i> spp.	0				▲				0
<i>Acinetobacter junii</i>	0				▲				0
Mycoplasmata	Result	-3	-2	-1	0	+1	+2	+3	Reference Interval
<i>Metamycoplasma hominis</i>	0				▲				0
Verrucomicrobiota	Result	-3	-2	-1	0	+1	+2	+3	Reference Interval
<i>Akkermansia muciniphila</i>	-1			▲					0



Microbiome Abundance Information:

- The GI360™ Microbiome Profile is a focused gut microbiota DNA analysis tool that identifies more than 45 targeted analytes across six phyla using a CE-marked multiplex PCR system. Patient results are compared to a highly defined normobiotic reference population (n > 1,100). The white shadowed web plot within the hexagonal diagram illustrates the degree to which an individual's microbiome profile deviates from normobiosis. The center of the diagram represents less bacterial abundance while the outer edges represent greater than normobiosis. Deviation from a hexagon-shaped plot indicates variant diversity of the microbial community. Key findings for patient's microbiome profile are summarized in the table below the diagram, and detailed results for all of the analytes are presented on the next 3 pages of the report. Detailed results for the specific bacteria are reported as -3 to +3 standard deviations, as compared to the normobiotic reference population.



Notes:

The gray-shaded area of the bar graph represents reference values outside the reporting limits for this test.

*This test was developed and its performance characteristics determined by Doctor's Data Laboratories in a manner consistent with CLIA requirements. The U. S. Food and Drug Administration (FDA) has not approved or cleared this test; however, FDA clearance is not currently required for clinical use. The results are not intended to be used as a sole means for clinical diagnosis or patient management decisions.

Methodology: Multiplex PCR



Order: 999999-9999



Client #: 999999

Doctor: Sample Doctor, MD
Doctors Data Inc
123 Main St.
St. Charles, IL 60174 USA

Patient: Sample Patient

Id: 999999

Age: 38 DOB: 00/00/1986

Sex: Female

Sample Collection

Date/Time

Date Collected

05/14/2025

Date Received

05/23/2025

Date Reported

06/02/2025

Specimens Collected 3

Pathogenic Bacteria	Result	NG	1+	2+	3+	4+	Reference Interval
<i>Aeromonas</i> spp.	NG	▲					No Growth
<i>Edwardsiella tarda</i>	NG	▲					No Growth
<i>Plesiomonas shigelloides</i>	NG	▲					No Growth
<i>Salmonella</i> group	NG	▲					No Growth
<i>Shigella</i> group	NG	▲					No Growth
<i>Vibrio cholerae</i>	NG	▲					No Growth
<i>Vibrio</i> spp.	NG	▲					No Growth
<i>Yersinia</i> spp.	NG	▲					No Growth
Imbalanced Bacteria	Result	NG	1+	2+	3+	4+	Reference Interval
<i>Bacillus licheniformis</i>	1+		▲				No Growth
Dysbiotic Bacteria	Result	NG	1+	2+	3+	4+	Reference Interval
<i>Citrobacter farmeri / amalonaticus</i>	4+					▲	No Growth
<i>Klebsiella oxytoca</i>	4+					▲	No Growth
<i>Klebsiella pneumoniae</i>	3+				▲		No Growth
<i>Providencia alcalifaciens</i>	4+					▲	No Growth
Yeast	Result	NG	1+	2+	3+	4+	Reference Interval
<i>Candida albicans</i>	1+		▲				0+ – 1+
<i>Candida krusei</i>	3+				▲		0+ – 1+
<i>Candida lusitanae</i>	1+		▲				0+ – 1+



Microbiology Information:

- Pathogenic bacteria** consist of known pathogenic bacteria that can cause disease in the GI tract. They are present due to the consumption of contaminated food or water, exposure to animals, fish, or amphibians known to harbor the organism. These organisms can be detected by either Multiplex PCR or microbiology culture.
- Imbalanced bacteria** are usually neither pathogenic nor beneficial to the host GI tract. Imbalances can occur when there are insufficient levels of beneficial bacteria and increased levels of commensal bacteria. Certain commensal bacteria are reported as dysbiotic at higher levels.
- Dysbiotic bacteria** consist of those bacteria that have the potential to cause disease in the GI tract. They can be present due to a number of factors including: exposure to chemicals that are toxic to beneficial bacteria; the use of antibiotics, oral contraceptives or other medications; poor fiber intake and high stress levels.

Notes:

NG = No Growth

Methodology: Culture and identification by MALDI-TOF and conventional biochemicals





Order: 999999-9999



Client #: 999999

Doctor: Sample Doctor, MD

Doctors Data Inc

123 Main St.

St. Charles, IL 60174 USA

Patient: Sample Patient

Id: 999999

Age: 38 **DOB:** 00/00/1986

Sex: Female

Sample Collection

Date/Time

Date Collected

05/14/2025

Date Received

05/23/2025

Date Reported

06/02/2025

Specimens Collected 3



Microbiology Information:

- **Yeast** may normally be present in small quantities on the skin, in the mouth and intestine. While small quantities of yeast may be normal, yeast observed in higher quantities is considered abnormal.





Order: 999999-9999



Client #: 999999

Doctor: Sample Doctor, MD
Doctors Data Inc
123 Main St.
St. Charles, IL 60174 USA

Patient: Sample Patient

Id: 999999

Age: 38 **DOB:** 00/00/1986

Sex: Female

Sample Collection

Date/Time

Date Collected

05/14/2025

Date Received

05/23/2025

Date Reported

06/02/2025

Specimens Collected 3

Viruses	Result	
Adenovirus F40/41	Negative	<input checked="" type="checkbox"/>
Norovirus GI/GII	Negative	<input checked="" type="checkbox"/>
Rotavirus A	Negative	<input checked="" type="checkbox"/>

Pathogenic Bacteria	Result	
<i>Campylobacter</i> (<i>C. jejuni</i> , <i>C. coli</i> and <i>C. lari</i>)	Negative	<input checked="" type="checkbox"/>
<i>Clostridioides difficile</i> (Toxin A/B)	Negative	<input checked="" type="checkbox"/>
<i>Escherichia coli</i> O157	Negative	<input checked="" type="checkbox"/>
Enterotoxigenic <i>Escherichia coli</i> (ETEC) lt/st	Negative	<input checked="" type="checkbox"/>
<i>Salmonella</i> spp.	Negative	<input checked="" type="checkbox"/>
Shiga-like toxin-producing <i>Escherichia coli</i> (STEC) stx1/stx2	Negative	<input checked="" type="checkbox"/>
<i>Shigella</i> (<i>S. boydii</i> , <i>S. sonnei</i> , <i>S. flexneri</i> & <i>S. dysenteriae</i>)	Negative	<input checked="" type="checkbox"/>
<i>Vibrio cholerae</i>	Negative	<input checked="" type="checkbox"/>

Parasites	Result	
<i>Cryptosporidium</i> (<i>C. parvum</i> and <i>C. hominis</i>)	Negative	<input checked="" type="checkbox"/>
<i>Entamoeba histolytica</i>	Negative	<input checked="" type="checkbox"/>
<i>Giardia duodenalis</i> (AKA <i>intestinalis</i> & <i>lamblia</i>)	Negative	<input checked="" type="checkbox"/>

Notes:

Methodology: Multiplex PCR





Order: 999999-9999



Client #: 999999

Doctor: Sample Doctor, MD
Doctors Data Inc
123 Main St.
St. Charles, IL 60174 USA

Patient: Sample Patient

Id: 999999

Age: 38 **DOB:** 00/00/1986

Sex: Female

Sample Collection

Date/Time

Date Collected

05/14/2025

Date Received

05/23/2025

Date Reported

06/02/2025

Specimens Collected

3

Protozoa

Result

<i>Balantidium coli</i>	Not Detected	<input checked="" type="checkbox"/>
<i>Blastocystis</i> spp.	Not Detected	<input checked="" type="checkbox"/>
<i>Chilomastix mesnili</i>	Not Detected	<input checked="" type="checkbox"/>
<i>Dientamoeba fragilis</i>	Not Detected	<input checked="" type="checkbox"/>
<i>Endolimax nana</i>	Not Detected	<input checked="" type="checkbox"/>
<i>Entamoeba coli</i>	Not Detected	<input checked="" type="checkbox"/>
<i>Entamoeba hartmanni</i>	Not Detected	<input checked="" type="checkbox"/>
<i>Entamoeba histolytica/Entamoeba dispar</i>	Not Detected	<input checked="" type="checkbox"/>
<i>Entamoeba polecki</i>	Not Detected	<input checked="" type="checkbox"/>
<i>Enteromonas hominis</i>	Not Detected	<input checked="" type="checkbox"/>
<i>Giardia duodenalis</i>	Not Detected	<input checked="" type="checkbox"/>
<i>Iodamoeba bütschlii</i>	Not Detected	<input checked="" type="checkbox"/>
<i>Isospora belli</i>	Not Detected	<input checked="" type="checkbox"/>
<i>Pentatrichomonas hominis</i>	Not Detected	<input checked="" type="checkbox"/>
<i>Retortamonas intestinalis</i>	Not Detected	<input checked="" type="checkbox"/>

Cestodes - Tapeworms

Result

<i>Diphyllobothrium latum</i>	Not Detected	<input checked="" type="checkbox"/>
<i>Dipylidium caninum</i>	Not Detected	<input checked="" type="checkbox"/>
<i>Hymenolepis diminuta</i>	Not Detected	<input checked="" type="checkbox"/>
<i>Hymenolepis nana</i>	Not Detected	<input checked="" type="checkbox"/>
<i>Taenia</i>	Not Detected	<input checked="" type="checkbox"/>

Trematodes - Flukes

Result

<i>Clonorchis sinensis</i>	Not Detected	<input checked="" type="checkbox"/>
<i>Fasciola hepatica/Fasciolopsis buski</i>	Not Detected	<input checked="" type="checkbox"/>
<i>Heterophyes heterophyes</i>	Not Detected	<input checked="" type="checkbox"/>
<i>Paragonimus westermani</i>	Not Detected	<input checked="" type="checkbox"/>

Nematodes - Roundworms

Result

<i>Ascaris lumbricoides</i>	Not Detected	<input checked="" type="checkbox"/>
-----------------------------	--------------	-------------------------------------

Notes:

Methodology: Microscopy



Order: 999999-9999



Client #: 999999

Doctor: Sample Doctor, MD
Doctors Data Inc
123 Main St.
St. Charles, IL 60174 USA

Patient: Sample Patient

Id: 999999

Age: 38 DOB: 00/00/1986

Sex: Female

Sample Collection

Date/Time

Date Collected

05/14/2025

Date Received

05/23/2025

Date Reported

06/02/2025

Specimens Collected 3

Nematodes - Roundworms

Result

<i>Capillaria hepatica</i>	Not Detected	<input checked="" type="checkbox"/>
<i>Capillaria philippinensis</i>	Not Detected	<input checked="" type="checkbox"/>
<i>Enterobius vermicularis</i>	Not Detected	<input checked="" type="checkbox"/>
Hookworm	Not Detected	<input checked="" type="checkbox"/>
<i>Strongyloides stercoralis</i>	Not Detected	<input checked="" type="checkbox"/>
<i>Trichuris trichiura</i>	Not Detected	<input checked="" type="checkbox"/>

Other Markers

Result

Reference Interval

Yeast	Moderate	<input checked="" type="checkbox"/>	Not Detected – Few
RBC	Not Detected	<input checked="" type="checkbox"/>	Not Detected – Few
WBC	Not Detected	<input checked="" type="checkbox"/>	Not Detected – Rare
Muscle fibers	Not Detected	<input checked="" type="checkbox"/>	Not Detected – Rare
Vegetable fibers	Rare	<input checked="" type="checkbox"/>	Not Detected – Few
Charcot-Leyden Crystals	Not Detected	<input checked="" type="checkbox"/>	Not Detected
Pollen	Not Detected	<input checked="" type="checkbox"/>	Not Detected
Mucus	Negative	<input checked="" type="checkbox"/>	Negative



Parasitology Information:

- This test is not designed to detect *Cyclospora cayetanensis* or *Microsporidia* spp.
- Intestinal parasites are abnormal inhabitants of the gastrointestinal tract that have the potential to cause damage to their host. The presence of any parasite within the intestine generally confirms that the patient has acquired the organism through fecal-oral contamination. Damage to the host includes parasitic burden, migration, blockage and pressure. Immunologic inflammation, hypersensitivity reactions and cytotoxicity also play a large role in the morbidity of these diseases. The infective dose often relates to severity of the disease and repeat encounters can be additive.
- There are two main classes of intestinal parasites, they include protozoa and helminths. The protozoa typically have two stages; the trophozoite stage that is the metabolically active, invasive stage and the cyst stage, which is the vegetative inactive form resistant to unfavorable environmental conditions outside the human host. Helminths are large, multicellular organisms. Like protozoa, helminths can be either free-living or parasitic in nature. In their adult form, helminths cannot multiply in humans.
- In general, acute manifestations of parasitic infection may involve diarrhea with or without mucus and or blood, fever, nausea, or abdominal pain. However these symptoms do not always occur. Consequently, parasitic infections may not be diagnosed or eradicated. If left untreated, chronic parasitic infections can cause damage to the intestinal lining and can be an unsuspected cause of illness and fatigue. Chronic parasitic infections can also be associated with increased intestinal permeability, irritable bowel syndrome, irregular bowel movements, malabsorption, gastritis or indigestion, skin disorders, joint pain, allergic reactions, and decreased immune function.

Notes:

Methodology: Microscopy, Macroscopic Observation



Order: 999999-9999



Client #: 999999

Doctor: Sample Doctor, MD

Doctors Data Inc

123 Main St.

St. Charles, IL 60174 USA

Patient: Sample Patient

Id: 999999

Age: 38 **DOB:** 00/00/1986

Sex: Female

Sample Collection

Date/Time

Date Collected

05/14/2025

Date Received

05/23/2025

Date Reported

06/02/2025


Specimens Collected

3



Parasitology Information:

- In some instances, parasites may enter the circulation and travel to various organs causing severe organ diseases such as liver abscesses and cysticercosis. In addition, some larval migration can cause pneumonia and in rare cases hyper infection syndrome with large numbers of larvae being produced and found in every tissue of the body.
- **Red Blood Cells** (RBC) in the stool may be associated with a parasitic or bacterial infection, or an inflammatory bowel condition such as ulcerative colitis. Colorectal cancer, anal fistulas, and hemorrhoids should also be ruled out.
- **White Blood Cells** (WBC) and **Mucus** in the stool can occur with bacterial and parasitic infections, with mucosal irritation, and inflammatory bowel diseases such as Crohn's disease or ulcerative colitis
- **Muscle fibers** in the stool are an indicator of incomplete digestion. Bloating, flatulence, feelings of "fullness" may be associated with increase in muscle fibers.
- **Vegetable fibers** in the stool may be indicative of inadequate chewing, or eating "on the run".





Order: 999999-9999



Client #: 999999

Doctor: Sample Doctor, MD
Doctors Data Inc
123 Main St.
St. Charles, IL 60174 USA

Patient: Sample Patient

Id: 999999

Age: 38 **DOB:** 00/00/1986

Sex: Female

Sample Collection

Date Collected

Date/Time

05/14/2025

Date Received

05/23/2025

Date Reported

06/02/2025

Specimens Collected

3

Digestion / Absorption	Result	Unit	L	WRI	H	Reference Interval
Elastase	>500	µg/g				> 200
Fat Stain	None					None – Moderate
Carbohydrates [†]	Negative					Negative
Inflammation	Result	Unit	L	WRI	H	Reference Interval
Lactoferrin	10.9	µg/mL				< 7.3
Lysozyme*	608	ng/mL				≤ 500
Calprotectin	133	µg/g				< 80
Immunology	Result	Unit	L	WRI	H	Reference Interval
Secretory IgA*	42.7	mg/dL				30 – 275
Short Chain Fatty Acids	Result	Unit	L	WRI	H	Reference Interval
% Acetate [‡]	84	%				50 – 72
% Propionate [‡]	6.2	%				11 – 25
% Butyrate [‡]	10	%				11 – 32
% Valerate [‡]	0.2	%				0.8 – 5.0
Butyrate [‡]	0.70	mg/mL				0.8 – 4.0
Total SCFA's [‡]	7.0	mg/mL				5.0 – 16.0
Intestinal Health Markers	Result	Unit	L	WRI	H	Reference Interval
pH	6.0					5.8 – 7.0
β-glucuronidase*	1790	U/h*g				2800 – 8000
Occult Blood	Negative					Negative
Macroscopic Appearance	Result	Unit				Reference Interval
Color	Brown					Brown
Consistency	Soft					Soft

Notes:

RI= Reference Interval, L (blue)= Low (below RI), WRI (green)= within RI, Yellow= moderately outside RI, L or H, H (red)= High (above RI)

*This test was developed and its performance characteristics determined by Doctor's Data Laboratories in a manner consistent with CLIA requirements. The U. S. Food and Drug Administration (FDA) has not approved or cleared this test; however, FDA clearance is not currently required for clinical use. The results are not intended to be used as a sole means for clinical diagnosis or patient management decisions.

[†]This test has been modified from the manufacturer's instructions and its performance characteristics determined by Doctor's Data Laboratories in a manner consistent with CLIA requirements.

[‡]This test was developed and its performance characteristics determined by Doctor's Data Laboratories in a manner consistent with CLIA requirements. The U.S. Food and Drug Administration (FDA) has not approved or cleared this test; however, FDA clearance is not currently required for clinical use.

Methodology: Turbidimetric immunoassay, Microscopy, Colormetric, Elisa, Gas Chromatography, Enzymatic, Guaiac, Macroscopic Observation



Order: 999999-9999



Client #: 999999

Doctor: Sample Doctor, MD

Doctors Data Inc

123 Main St.

St. Charles, IL 60174 USA

Patient: Sample Patient

Id: 999999

Age: 38 **DOB:** 00/00/1986

Sex: Female

Sample Collection

Date/Time

Date Collected

05/14/2025

Date Received

05/23/2025

Date Reported

06/02/2025

Specimens Collected

3



Chemistry Information:

- **Elastase** findings can be used for assessing pancreatic exocrine function and insufficiency.
- **Fat Stain:** Microscopic determination of fecal fat using Sudan IV staining is a qualitative procedure utilized to assess fat absorption and to detect steatorrhea.
- **Carbohydrates:** The presence of reducing substances in stool specimens can indicate carbohydrate malabsorption.
- **Lactoferrin** and **Calprotectin** are reliable markers for differentiating organic inflammation (IBD) from function symptoms (IBS) and for management of IBD. Monitoring levels of fecal lactoferrin and calprotectin can play an essential role in determining the effectiveness of therapy, are good predictors of IBD remission, and can indicate a low risk of relapse.
- **Lysozyme** is an enzyme secreted at the site of inflammation in the GI tract and elevated levels have been identified in IBD patients.
- **Secretory IgA** (sIgA) is secreted by mucosal tissue and represents the first line of defense of the GI mucosa and is central to the normal function of the GI tract as an immune barrier. Elevated levels of sIgA have been associated with an upregulated immune response.
- **Short chain fatty acids (SCFAs):** SCFAs are the end product of the bacterial fermentation process of dietary fiber by beneficial flora in the gut and play an important role in the health of the GI as well as protecting against intestinal dysbiosis. Lactobacilli and bifidobacteria produce large amounts of short chain fatty acids, which decrease the pH of the intestines and therefore make the environment unsuitable for pathogens, including bacteria and yeast. Studies have shown that SCFAs have numerous implications in maintaining gut physiology. SCFAs decrease inflammation, stimulate healing, and contribute to normal cell metabolism and differentiation. Levels of **Butyrate** and **Total SCFA** in mg/mL are important for assessing overall SCFA production, and are reflective of beneficial flora levels and/or adequate fiber intake.
- **pH:** Fecal pH is largely dependent on the fermentation of fiber by the beneficial flora of the gut.
- **Occult blood:** A positive occult blood indicates the presence of free hemoglobin found in the stool, which is released when red blood cells are lysed.
- **β -glucuronidase** is an enzyme that breaks the tight bond between glucuronic acid and toxins in the intestines. The binding of toxins in the gut is protective by way of blocking their absorption and facilitating excretion.





Order: 999999-9999



Client #: 999999

Doctor: Sample Doctor, MD
Doctors Data Inc
123 Main St.
St. Charles, IL 60174 USA

Patient: Sample Patient

Id: 999999

Age: 38 **DOB:** 00/00/1986

Sex: Female

Sample Collection

Date/Time

Date Collected

05/14/2025

Date Received

05/23/2025

Date Reported


06/02/2025

Specimens Collected


3

Citrobacter farmeri / amalonaticus

Natural Agents



Prescriptive Agents



Susceptibility Information:

- Natural antibacterial** agents may be useful for treatment of patients when organisms display in-vitro sensitivity to these agents. The test is performed by using standardized techniques and filter paper disks impregnated with the listed agent. Relative sensitivity is reported for each natural agent based upon the diameter of the zone of inhibition surrounding the disk. Data based on over 5000 individual observations were used to relate the zone size to the activity level of the agent. A scale of relative sensitivity is defined for the natural agents tested.
- Susceptible** results imply that an infection due to the bacteria may be appropriately treated when the recommended dosage of the tested antimicrobial agent is used. **Intermediate** results imply that response rates may be lower than for susceptible bacteria when the tested antimicrobial agent is used. **Resistant** results imply that the bacteria will not be inhibited by normal dosage levels of the tested antimicrobial agent.

Notes:

*This test was developed and its performance characteristics determined by Doctor's Data Laboratories in a manner consistent with CLIA requirements. The U. S. Food and Drug Administration (FDA) has not approved or cleared this test; however, FDA clearance is not currently required for clinical use. The results are not intended to be used as a sole means for clinical diagnosis or patient management decisions.





Order: 999999-9999



Client #: 999999

Doctor: Sample Doctor, MD

Doctors Data Inc

123 Main St.

St. Charles, IL 60174 USA

Patient: Sample Patient

Id: 999999

Age: 38 **DOB:** 00/00/1986

Sex: Female

Sample Collection

Date/Time

Date Collected

05/14/2025

Date Received

05/23/2025

Date Reported


06/02/2025

Specimens Collected


3

Klebsiella oxytoca

Natural Agents



Prescriptive Agents



Susceptibility Information:

- Natural antibacterial** agents may be useful for treatment of patients when organisms display in-vitro sensitivity to these agents. The test is performed by using standardized techniques and filter paper disks impregnated with the listed agent. Relative sensitivity is reported for each natural agent based upon the diameter of the zone of inhibition surrounding the disk. Data based on over 5000 individual observations were used to relate the zone size to the activity level of the agent. A scale of relative sensitivity is defined for the natural agents tested.
- Susceptible** results imply that an infection due to the bacteria may be appropriately treated when the recommended dosage of the tested antimicrobial agent is used. **Intermediate** results imply that response rates may be lower than for susceptible bacteria when the tested antimicrobial agent is used. **Resistant** results imply that the bacteria will not be inhibited by normal dosage levels of the tested antimicrobial agent.

Notes:

*This test was developed and its performance characteristics determined by Doctor's Data Laboratories in a manner consistent with CLIA requirements. The U. S. Food and Drug Administration (FDA) has not approved or cleared this test; however, FDA clearance is not currently required for clinical use. The results are not intended to be used as a sole means for clinical diagnosis or patient management decisions.





Order: 999999-9999



Client #: 999999

Doctor: Sample Doctor, MD
Doctors Data Inc
123 Main St.
St. Charles, IL 60174 USA

Patient: Sample Patient

Id: 999999

Age: 38 DOB: 00/00/1986

Sex: Female

Sample Collection

Date/Time

Date Collected

05/14/2025

Date Received

05/23/2025

Date Reported

06/02/2025

Specimens Collected

3

Klebsiella pneumoniae

Natural Agents

	Low Inhibition	High Inhibition
Caprylic Acid*		▲
Uva Ursi*	▲	
Olive Leaf Extract*	▲	
Oregano*	▲	
Goldenseal*	▲	
Ionic Silver*		▲
Colloidal Silver*		▲

Prescriptive Agents

	Resistant	Intermediate	Susceptible
Amoxicillin-Clavulanic Acid			<input checked="" type="checkbox"/>
Ampicillin	<input checked="" type="checkbox"/>		
Cefazolin			<input checked="" type="checkbox"/>
Ceftazidime			<input checked="" type="checkbox"/>
Ciprofloxacin			<input checked="" type="checkbox"/>
Sulfamethoxazole / Trimethoprim			<input checked="" type="checkbox"/>



Susceptibility Information:

- Natural antibacterial** agents may be useful for treatment of patients when organisms display in-vitro sensitivity to these agents. The test is performed by using standardized techniques and filter paper disks impregnated with the listed agent. Relative sensitivity is reported for each natural agent based upon the diameter of the zone of inhibition surrounding the disk. Data based on over 5000 individual observations were used to relate the zone size to the activity level of the agent. A scale of relative sensitivity is defined for the natural agents tested.
- Susceptible** results imply that an infection due to the bacteria may be appropriately treated when the recommended dosage of the tested antimicrobial agent is used. **Intermediate** results imply that response rates may be lower than for susceptible bacteria when the tested antimicrobial agent is used. **Resistant** results imply that the bacteria will not be inhibited by normal dosage levels of the tested antimicrobial agent.

Notes:

*This test was developed and its performance characteristics determined by Doctor's Data Laboratories in a manner consistent with CLIA requirements. The U. S. Food and Drug Administration (FDA) has not approved or cleared this test; however, FDA clearance is not currently required for clinical use. The results are not intended to be used as a sole means for clinical diagnosis or patient management decisions.





Order: 999999-9999



Client #: 999999

Doctor: Sample Doctor, MD
Doctors Data Inc
123 Main St.
St. Charles, IL 60174 USA

Patient: Sample Patient

Id: 999999

Age: 38 **DOB:** 00/00/1986

Sex: Female

Sample Collection

Date/Time

Date Collected

05/14/2025

Date Received

05/23/2025

Date Reported


06/02/2025

Specimens Collected


3

Providencia alcalifaciens

Natural Agents



Prescriptive Agents



Susceptibility Information:

- Natural antibacterial** agents may be useful for treatment of patients when organisms display in-vitro sensitivity to these agents. The test is performed by using standardized techniques and filter paper disks impregnated with the listed agent. Relative sensitivity is reported for each natural agent based upon the diameter of the zone of inhibition surrounding the disk. Data based on over 5000 individual observations were used to relate the zone size to the activity level of the agent. A scale of relative sensitivity is defined for the natural agents tested.
- Susceptible** results imply that an infection due to the bacteria may be appropriately treated when the recommended dosage of the tested antimicrobial agent is used. **Intermediate** results imply that response rates may be lower than for susceptible bacteria when the tested antimicrobial agent is used. **Resistant** results imply that the bacteria will not be inhibited by normal dosage levels of the tested antimicrobial agent.

Notes:

*This test was developed and its performance characteristics determined by Doctor's Data Laboratories in a manner consistent with CLIA requirements. The U. S. Food and Drug Administration (FDA) has not approved or cleared this test; however, FDA clearance is not currently required for clinical use. The results are not intended to be used as a sole means for clinical diagnosis or patient management decisions.





Order: 999999-9999



Client #: 999999

Doctor: Sample Doctor, MD
Doctors Data Inc
123 Main St.
St. Charles, IL 60174 USA

Patient: Sample Patient

Id: 999999

Age: 38 DOB: 00/00/1986

Sex: Female

Sample Collection

Date/Time

Date Collected

05/14/2025

Date Received




05/23/2025

Date Reported

06/02/2025

Specimens Collected

3

Candida krusei**Natural Agents****Non-Absorbed Antifungals****Azole Antifungals****Susceptibility Information:**

- Natural antifungal** agents may be useful for treatment of patients when organisms display in-vitro susceptibility to these agents. The test is performed by using standardized techniques and filter paper disks impregnated with the listed agent. Relative activity is reported for each natural agent based upon the diameter of the zone of inhibition or no growth zone surrounding the disk. Data based on over 5000 individual observations were used to relate the zone size to the activity level of the agent. A scale of relative activity is defined for the natural agents tested.
- Non-absorbed antifungals** may be useful for treatment of patients when organisms display in-vitro susceptibility to these agents. The test is performed using standardized commercially prepared disks impregnated with Nystatin. Relative activity is reported based upon the diameter of the zone of inhibition or no growth zone surrounding the disk.
- Susceptible** results imply that an infection due to the fungus may be appropriately treated when the recommended dosage of the tested antifungal agent is used. **Susceptible - Dose Dependent (S-DD)** results imply that an infection due to the fungus may be treated when the highest recommended dosage of the tested antifungal agent is used. **Resistant** results imply that the fungus will not be inhibited by normal dosage levels of the tested antifungal agent.

Notes:

*This test was developed and its performance characteristics determined by Doctor's Data Laboratories in a manner consistent with CLIA requirements. The U. S. Food and Drug Administration (FDA) has not approved or cleared this test; however, FDA clearance is not currently required for clinical use. The results are not intended to be used as a sole means for clinical diagnosis or patient management decisions.





Order: 999999-9999



Client #: 999999

Doctor: Sample Doctor, MD

Doctors Data Inc

123 Main St.

St. Charles, IL 60174 USA

Patient: Sample Patient

Id: 999999

Age: 38 DOB: 00/00/1986

Sex: Female

Sample Collection

Date/Time

Date Collected

05/14/2025

Date Received

05/23/2025

Date Reported

06/02/2025

Specimens Collected 3

Introduction

This analysis of the stool specimen provides fundamental information about the overall gastrointestinal health of the patient. When abnormal microflora or significant aberrations in intestinal health markers are detected, specific commentaries are presented. If no significant abnormalities are found, commentaries are not presented.

The majority of reference intervals are established from adult populations. Results may differ in pediatric populations and care should be taken when interpreting these values.

Microbiome Abundance Information

Actinobacteria (phylum)

Actinobacteria is one of the largest bacterial phyla, comprised of Gram-positive bacteria. This phylum includes a wide range of species, with different morphological and physiological characteristics. Significant groups in the human colon include Actinomycetales and Bifidobacteriales. Actinomycetales were inversely associated with clinically significant depression in IBS patients, suggesting these bacteria may be depleted in depressed IBS patients. A strict vegetarian diet may increase the total count of *Actinomyces* spp. compared to following a Western diet.

↓ Actinomycetales (order)

Actinomycetales are considered low abundance colonizers of the gastrointestinal tract with primary residence on the skin. Intake of proton-pump inhibitor drugs has been shown to increase the abundance of Actinomycetales in the gut, possibly by reducing gastric acidity and enabling intestinal colonization by oral microbes. Actinomycetales may be depleted in depressed irritable bowel syndrome patients. The abundance of *Actinomyces* spp. was shown to be higher with a strict vegetarian diet compared to a common Western diet.

↓ Bifidobacterium (genus)

Considered amongst the most beneficial commensal bacteria in the human gut, *Bifidobacterium* spp. are able to degrade monosaccharides, galacto-, manno-, and fructo-oligosaccharides, as well as some complex carbohydrates. Many of the non-digestible oligosaccharides, found as natural components in mother's milk, select for colonization of these species which dominate the infant gut shortly after birth. Bifidobacteria may provide health benefits directly through interactions with the host, and indirectly through interactions with other microorganisms. *Bifidobacterium* spp. take part in production and adsorption of vitamins, such as vitamins K and B12, biotin, folate, thiamine, riboflavin, and pyridoxine. They are also involved in lipid absorption and metabolism, glucose and energy homeostasis, and regulating intestinal barrier function. Although *Bifidobacterium* produce acetate over butyrate, healthy levels of *Bifidobacterium* spp. facilitate colonization of *Faecalibacterium prausnitzii*. Polyphenols derived from chocolate, green tea, blackcurrant, red wine and grape seed extracts have been shown to increase *Bifidobacterium* species. The increased abundance of *Bifidobacterium* species has been associated with amelioration of inflammation. Multiple published studies have suggested that there is an association between obesity and a lower abundance of bifidobacteria. They may also be less abundant in elderly populations, patients with rheumatoid arthritis, and in individuals diagnosed with Alzheimer's disease. Patients with active inflammatory bowel disease (IBD) have a lower abundance of *Bifidobacterium* spp. than patients whose IBD is in remission. Taking a probiotic containing bifidobacteria, lactobacilli, and streptococci might help in controlling ulcerative colitis symptoms and preventing their recurrence. Some *Bifidobacterium* strains have been shown to have beneficial effects in irritable bowel syndrome (IBS). *Bifidobacterium* spp. abundance has been shown to be diminished with IBD and with long term use of macrolide antibiotics. Luminal bifidobacteria is reduced with restriction of fermentable carbohydrates, i.e. a low FODMAP diet. High fat dietary feeding is also associated with reduced abundance of bifidobacteria. Consumption of maize and barley-based whole grain products and red berries, which are comprised of anthocyanins, are known to increase levels of bifidobacteria.

Bacteroidetes (phylum)

Bacteroidetes make up approximately 28% of the gut microbiota in healthy human adults. They are early colonizers of the infant gut and are amongst the most stable, at a species and strain level, in the host. A low preponderance of Bacteroidetes in relation to Firmicutes has been associated with obesity, though this can increase with weight loss and restricted calorie intake.



Order: 999999-9999



Client #: 999999

Doctor: Sample Doctor, MD

Doctors Data Inc

123 Main St.

St. Charles, IL 60174 USA

Patient: Sample Patient

Id: 999999

Age: 38 DOB: 00/00/1986

Sex: Female

Sample Collection

Date/Time

Date Collected

05/14/2025

Date Received

05/23/2025

Date Reported

06/02/2025

Specimens Collected 3

Microbiome Abundance Information continued...

↓ *Alistipes* (genus)

Alistipes does not contribute significantly to short chain fatty acid production. A diet rich in animal protein and fat increases the abundance of *Alistipes*. High abundance of *Alistipes* was identified as a possible predictor of successful weight loss. Increased abundance of *Alistipes* has been correlated with a greater frequency of pain in pediatric irritable bowel syndrome patients. In contrast, *Alistipes onderdonkii* was shown to be decreased in patients diagnosed with ulcerative colitis. Lower abundance of the *Alistipes* genus has been observed in patients with psoriatic arthritis and pediatric Crohn's disease. *Alistipes* may positively correlate with depression.

↓ *Bacteroides pectinophilus* (species)

Bacteroides pectinophilus contributes to breakdown of dietary pectins which are prebiotics. Pectins are complex, plant-derived carbohydrates that are indigestible by human enzymes, but can be easily degraded by certain commensal bacteria in the gut. Subsequent microbial fermentation of constituent sugar moieties yields important short chain fatty acids and other metabolites. The pectin-derived microbial fermentation products have important functions including reduction of ammonia, delay of gastric emptying and postprandial glucose regulation, induction of gut immunity, and maintenance of the mucosal barrier. Adequate intake and microbial metabolism of pectins appears to stimulate growth of various beneficial bacteria, including *Lachnospiraceae*, *Dorea* species, *Bifidobacterium*, *Lactobacillus* species, *Faecalibacterium prausnitzii*, and *Eubacterium rectale*. The abundance of *B. pectinophilus* has been positively correlated with a healthy fasting serum lipid profile, and negatively correlated with biomarkers of for insulin resistance and dyslipidemia. *B. pectinophilus* was less abundant for IBS patients compared to healthy controls. High consumption of kimchi (fermented cabbage) may be associated with lower than normal levels of *B. pectinophilus*.

↓ *Prevotella* (genus)

Prevotella-rich dysbiosis has been associated with insulin-resistance, obesity and hypertension. *Prevotella* have been shown to be significantly decreased in Crohn's disease and Parkinson's disease. High levels of fiber and carbohydrates from fruits and vegetables in a Mediterranean diet have been shown to increase the relative abundance of *Prevotella*.

↓ *Bacteroides* (species)

Species in the genus *Bacteroides* carry out broad metabolic functions, including degradation of complex plant polysaccharides, proteolytic activities, de-conjugation of bile acids, mucosal barrier integrity, short chain fatty acid production, fatty acid storage and glucose metabolism. *Bacteroides* spp. are maintained at a higher abundance in breastfed individuals into adulthood. *Bacteroides fragilis* plays an important role in the prevention of intestinal inflammation. An energy-restricted diet has been shown to increase *B. fragilis* in overweight adolescents. An increase in *B. stercoris* has been associated with higher risk of colon cancer. Decreased levels of *Bacteroides* spp. have been reported in association with multiple sclerosis, rheumatoid arthritis and Parkinson's disease.

↓ *Parabacteroides* (genus)

The abundance of *Parabacteroides* spp., major anaerobic producers of acetate and succinate is increased with a high fat diet and is positively correlated with body weight. *Parabacteroides* spp., along with certain *Bacteroides* spp., have been shown to distinguish healthy adults from patients with irritable bowel syndrome or ulcerative colitis. Reduced abundance of this group of bacteria has also been linked to Crohn's disease in children. *Parabacteroides* spp. has been found to be less abundant in patients with multiple sclerosis.

Firmicutes (phylum)

The phylum Firmicutes constitutes the most diverse and abundant group of gastrointestinal microbiota which are grouped into four classes, Bacilli, Clostridia, Erysipelotrichia, and Negativicutes. They constitute about 39% of gut bacteria in healthy adults, but may increase to as high as 80% in an imbalanced microbial community.

↑ *Clostridium* (genus)

Clostridium spp. represents an extremely heterogeneous class of organisms that are still actively undergoing taxonomic revision. *Clostridium* spp. are strict anaerobic, spore-forming bacteria. Decreased abundance of the genus *Clostridium* was found to be associated with prediabetes. Some *Clostridium* spp. are transferred to infants from breast milk within the first months of life. Increased levels of some *Clostridium* spp. were observed in irritable bowel syndrome patients. Many species, some of them related to diarrhea, were decreased after consumption of inulin combined with maltodextrin.



Order: 999999-9999



Client #: 999999

Doctor: Sample Doctor, MD

Doctors Data Inc

123 Main St.

St. Charles, IL 60174 USA

Patient: Sample Patient

Id: 999999

Age: 38 DOB: 00/00/1986

Sex: Female

Sample Collection

Date/Time

Date Collected

05/14/2025

Date Received

05/23/2025

Date Reported

06/02/2025

Specimens Collected 3

Microbiome Abundance Information continued...

↓ *Clostridium methylpentosum* (species)

Appropriate digestion and metabolism of complex dietary carbohydrates from plants drives healthy diversity in the gut microbiota. *Clostridium methylpentosum* ferments the naturally occurring sugar L-rhamnose that is released by microbial breakdown of plant-derived pectin. Rhamnose is fermented to propionate and acetate which are short chain fatty acids that have pivotal regulatory roles in the maintenance of mucosal barrier integrity, gut microbiota balance, post-prandial appetite suppression and normoglycemia. Lower levels of *C. methylpentosum* were reported for children with autism and pervasive developmental disorder compared to neurotypicals controls. Consumption of probiotic yogurt LKM512 containing *Bifidobacterium animalis* (subspecies lactis LKM512) increased the levels of *C. methylpentosum*.

↓ *Dorea* (genus)

Dorea is a genus within the *Lachnospiraceae* family that is in the Firmicutes phylum. *Dorea* species are known to produce hydrogen and carbon dioxide as end-products of glucose fermentation and may be associated with bloating. Decreased levels of *Dorea* spp. were observed in patients with Parkinson's disease. Recent studies have identified increased levels of *Dorea* spp. in patients diagnosed with IBS, nonalcoholic fatty liver disease and non-alcoholic steatohepatitis, multiple sclerosis and colorectal cancer.

↓ *Anaerobutyricum hallii* (species)

Anaerobutyricum hallii and *Agathobacter rectalis* (*Eubacterium rectale*) are both part of the *Lachnospiraceae* family that is in the Firmicutes family. *A. hallii* and *A. rectalis* produce butyrate that is a key regulator of mucosal barrier integrity and function. Decreased levels of *Anaerobutyricum*/*Agathobacter* spp have been associated with very high protein diets. *Anaerobutyricum hallii* is capable of metabolizing glucose products with antimicrobial properties.

↓ *Agathobacter rectalis* (*Eubacterium rectale*)

Agathobacter rectalis (*Eubacterium rectale*) is part of the *Lachnospiraceae* family and produce butyrate. *Agathobacter rectalis* was found to be in lower abundance in patients with type 2 diabetes, colorectal cancer, and chronic idiopathic diarrhea. There is a negative correlation between *Agathobacter rectalis* and the symptomology of irritable bowel syndrome (IBS). Decreased levels of *Anaerobutyricum*/*Agathobacter* spp. have been associated with very high protein diets.

↓ *Faecalibacterium prausnitzii* (species)

Faecalibacterium prausnitzii is one of the most abundant butyrate producing bacteria in a healthy gastrointestinal tract. As such, *F. prausnitzii* is a protective factor for the intestinal mucosa and supports very important intestinal barrier functions. *F. prausnitzii* exerts anti-inflammatory effects via metabolites such as short-chain fatty acids. *F. prausnitzii* is reduced in inflammatory bowel disease, irritable bowel syndrome, celiac disease and gastrointestinal inflammation in general. It is reduced in patients diagnosed with Parkinson's disease, bipolar disorder, colorectal cancer, diabetes and chronic idiopathic diarrhea. Diminished levels of *F. prausnitzii* were found in patients with major depressive disorder. The abundance of *F. prausnitzii* together with *E. coli* has been proposed as a discrimination tool between ulcerative colitis and Crohn's disease. *F. prausnitzii* has been correlated with pediatric obesity in instances of high consumption of foods that are rich in unabsorbed carbohydrate (banana, maize, rice). The prebiotic inulin has been shown to increase the proportion of *F. prausnitzii* in the human intestinal microbiota. Low FODMAP diets are associated with diminished *F. prausnitzii* and butyrate production.

↑ *Lachnospiraceae* (family)

The *Lachnospiraceae* family is a diverse group of butyric acid producers, which have been associated with beneficial microbial and epithelial cell growth. Consumption of a Mediterranean diet decreased levels of species belonging to *Lachnospiraceae*. *Lachnospiraceae* are known to increase with intake of cruciferous vegetables and wheat bran, and decrease with a resistant starch diet.



Order: 999999-9999



Client #: 999999

Doctor: Sample Doctor, MD
Doctors Data Inc
123 Main St.
St. Charles, IL 60174 USA

Patient: Sample Patient

Id: 999999

Age: 38 DOB: 00/00/1986

Sex: Female

Sample Collection

Date/Time

Date Collected

05/14/2025

Date Received

05/23/2025

Date Reported

06/02/2025

Specimens Collected

3

Microbiome Abundance Information continued...

↑ *Ruminococcus/Mediterraneibacter* (genus)

Members of the *Ruminococcus* and the new genus *Mediterraneibacter* sensu produce acetate, but not butyrate. *Mediterraneibacter* (*Ruminococcus*) *gnavus*, like *Akkermansia muciniphila* is a mucin degrading specialist. Higher levels of *Ruminococcus/Mediterraneibacter* were associated with non-alcoholic fatty liver disease and non-alcoholic steatohepatitis. Reduced levels of *Ruminococcus bromii* were observed in patients with primary biliary cirrhosis. Increased abundance of *Ruminococcus/Mediterraneibacter* spp. has been reported in irritable bowel syndrome (IBS), whereas *Ruminococcus/Mediterraneibacter* spp. are reportedly decreased in abundance with Crohn's disease and ulcerative colitis. *Mediterraneibacter gnavus* has been found to be in higher abundance in diarrhea predominant IBS. Intake of resistant starch has been associated with increased levels of *R. bromii*, whereas a diet rich in animal protein and fat was found to reduce the abundance of this species in the human gut.

↑ *Streptococcus* (genus)

Higher abundance of *S. salivarius* and *S. thermophilus* (Firmicutes phylum) have been associated with a moderate to severe disease course in newly diagnosed ulcerative colitis (UC) patients. These findings are in accordance with a study that showed that UC patients have significantly increased *Streptococcus* spp. and depletion of *Bifidobacterium* spp. Higher levels of *Streptococcus* spp. were also observed in patients with colorectal cancer compared to healthy controls. Administration of *S. salivarius* together with *Bifidobacterium bifidum* was shown to reduce the incidence of acute diarrhea and rotavirus shedding in infants. *S. salivarius* and *S. thermophilus* are also widely used in dairy products like yogurt and cheese.

Proteobacteria (phylum)

Proteobacteria include a wide variety of pathogens, including species within the *Escherichia*, *Shigella*, *Salmonella*, *Vibrio*, and *Helicobacter* genera. The phylum includes a number of species that are permanent residents of the microbiota and capable of inducing nonspecific inflammation and diarrhea when their presence is increased. Proteobacteria make up approximately 2% of the gut microbiota in healthy adults.

↑ Proteobacteria

A high-fat diet is positively associated with an abundance of Proteobacteria. Slightly increased abundance of Proteobacteria may be associated with low-grade inflammation. Proteobacteria are increased in inflammatory bowel disease and irritable bowel syndrome. Higher abundance of Proteobacteria has been associated with a moderate to severe disease course in newly discovered ulcerative colitis patients. They are associated with diarrhea in IBS.

↑ *Enterobacteriaceae* (family)

Enterobacteriaceae is a large family of bacteria within the Proteobacteria phyla. *Enterobacteriaceae* is inclusive of normal commensal species, harmless opportunists, and many of the more familiar pathogens, such as *Salmonella*, *Escherichia coli*, *Klebsiella*, *Shigella* and *Proteus*. Other potential disease-causing bacteria in this family include *Enterobacter* and *Citrobacter* species. The abundance of Proteobacteria, which are generally pro-inflammatory, is presented on the white shadowed web plot within the hexagonal diagram. The presence of specific dysbiotic and pathogenic *Enterobacteriaceae* bacteria, if detected by PCR or culture, are reported in the Gastrointestinal Pathogens and Microbiology sections of this report.

Overall, *Enterobacteriaceae* were found at higher levels in patients with NAFLD and PD. Diets rich in in complex carbohydrates are associated with lower levels of *Enterobacteriaceae*, in comparison to diets rich in fat and/or protein.

Mycoplasmata (Tenericutes) (phylum)

Mycoplasmata are cell wall-less bacteria that do not synthesize precursors of peptidoglycan. Mycoplasmata consist of four main clades designated as the *Acholeplasma*, *Spiroplasma*, *Pneumoniae* and *Hominis* clusters. Mycoplasmatas are typically parasites or commensals of eukaryotic hosts.

Verrucomicrobiota (Verrucomicrobia) (phylum)

Verrucomicrobiota is a less common phylum in the human microbiota, but one with increasing recognition with regards to health. Verrucomicrobiota includes *Akkermansia muciniphila*. The obligate anaerobe *A. muciniphila* constitutes 3-5% of total bacteria in a healthy microbiome, and has a protective or anti-inflammatory role in the intestinal mucosa.



Order: 999999-9999



Client #: 999999

Doctor: Sample Doctor, MD

Doctors Data Inc

123 Main St.

St. Charles, IL 60174 USA

Patient: Sample Patient

Id: 999999

Age: 38 DOB: 00/00/1986

Sex: Female

Sample Collection

Date/Time

Date Collected

05/14/2025

Date Received

05/23/2025

Date Reported

06/02/2025

Specimens Collected 3

Microbiome Abundance Information continued...

↓ *Akkermansia muciniphila* (genus)

Higher abundance of *Akkermansia muciniphila* has been associated with a milder disease course in newly discovered ulcerative colitis patients. Archaea and *Akkermansia* were significantly more prevalent after weight reduction. A Low FODMAP diet has been shown to decrease the abundance of *A. muciniphila* leading to recommendations against long-term use of such a diet. *A. muciniphila* is a mucolytic specialist that has potent anti-inflammatory effects in part associated with a specific surface coat protein (Amuc- 1100).

Microbiology

Pathogenic/Dysbiotic Flora

In a healthy balanced state of intestinal flora, the beneficial bacteria make up a significant proportion of the total microflora. However, in many individuals there is an imbalance or deficiency of beneficial flora (insufficiency dysbiosis) and an overgrowth of non-beneficial (imbalance) or even pathogenic microorganisms. This can be due to a number of factors including: consumption of contaminated water or food; daily exposure of chemicals that are toxic to beneficial bacteria; the use of antibiotics, oral contraceptives or other medications; poor fiber intake and high stress levels.

A number of toxic substances can be produced by the dysbiotic bacteria including amines, ammonia, hydrogen sulfide, phenols, and secondary bile acids which may cause inflammation or damage to the brush border of the intestinal lining. If left unchecked, long-term damage to the intestinal lining may result in leaky gut syndrome, allergies, autoimmune disease (e.g. rheumatoid arthritis), irritable bowel syndrome, fatigue, chronic headaches, and sensitivities to a variety of foods. In addition, pathogenic bacteria can cause acute symptoms such as abdominal pain, nausea, diarrhea, vomiting, and fever in cases of food poisoning.

Bacterial sensitivities to a variety of prescriptive and natural agents have been provided for the pathogenic bacteria that were cultured from this patient's specimen. This provides the practitioner with useful information to help plan an appropriate treatment regimen. Supplementation with probiotics or consumption of foods (yogurt, kefir, miso, tempeh, tamari sauce) containing strains of lactobacilli, bifidobacteria, and enterococci may help restore healthy flora levels. Soluble fiber and polyphenols derived from chocolate, green tea, blackcurrant, red wine and grape seed extracts have been found to increase the numbers of beneficial bacteria. Hypochlorhydria may also predispose an individual to bacterial overgrowth, particularly in the small intestine. Nutritional anti-inflammatories can aid in reversing irritation to the GI lining. These include quercetin, vitamin C, curcumin, gamma-linoleic acid, omega-3 fatty acids (EPA, DHA), and aloe vera. Other nutrients such as zinc, beta-carotene, pantothenic acid, and L-glutamine provide support for regeneration of the GI mucosa. A comprehensive program may be helpful in individuals in whom a dysbiotic condition has caused extensive GI damage.

Citrobacter spp

Citrobacter spp., a gram-negative bacterium and member of the *Enterobacteriaceae* family, is considered dysbiotic at 3+ or greater. *Citrobacter freundii* complex (including *C. freundii*, *C. braakii*, *C. gullenii*, *C. murlinae*, *rodentium*, *C. wermanii*, *C. youngae*), *C. koseri* and *C. farmeri*, can cause diarrheal disease. Symptoms are the result of an *E. coli*-like heat-stable enterotoxin and hydrogen sulfide. *Citrobacter freundii* complex has been implicated as a cause of gastrointestinal infection and inflammation, acute dysentery, and dyspepsia. Acute symptoms can include profuse, watery diarrhea without abdominal pain, fecal blood, or white blood cells.

Citrobacter spp. thrive on fructooligosaccharides (FOS), a common ingredient in artificial or alternative sweetener.

Antibiotics may be indicated if symptoms are prolonged. Refer to the antimicrobial susceptibilities to identify the most appropriate agent.



Order: 999999-9999



Client #: 999999

Doctor: Sample Doctor, MD

Doctors Data Inc

123 Main St.

St. Charles, IL 60174 USA

Patient: Sample Patient

Id: 999999

Age: 38 DOB: 00/00/1986

Sex: Female

Sample Collection

Date/Time

Date Collected

05/14/2025

Date Received

05/23/2025

Date Reported

06/02/2025

Specimens Collected 3

Microbiology continued...

Klebsiella spp

Klebsiella spp. are gram-negative bacilli belonging to the *Enterobacteriaceae* family and closely related to the genera *Enterobacter* and *Serratia*. *Klebsiella* spp. are considered dysbiotic in the amount of 3 - 4 +. *Klebsiella* spp. are widely distributed in nature and in the gastrointestinal tract of humans. In humans, they may colonize the skin, oral cavity, pharynx, or gastrointestinal tract. Regarded as normal flora in many parts of the colon, intestinal tract and biliary tract, the gut is the main reservoir of opportunistic strains. This bacteria has the potential to cause intestinal, lung, urinary tract, and wound infections, but overgrowth of *Klebsiella* spp. is commonly asymptomatic. *K. pneumoniae*, in particular, may cause diarrhea and some strains are enterotoxigenic. Infection has been linked to ankylosing spondylitis as well as myasthenia gravis (antigenic cross-reactivity), and these patients usually carry larger numbers of the organism in their intestines than healthy individuals. *Klebsiella oxytoca* causes antibiotic associated hemorrhagic colitis. These strains have been shown to produce a cytotoxin that is capable of inducing cell death in various epithelial-cell cultures.

Klebsiella is a significant nosocomial infectious agent, partially due to the ability of organisms to spread rapidly. *Klebsiella* accounts for approximately 3-7% of all hospital-acquired infections, placing it among the top eight pathogens in hospitals. Extraintestinal infection typically involves the respiratory or urinary tracts, but may infect other areas such as the biliary tract and surgical wound sites. *K. pneumoniae* and *K. oxytoca* are the two members of this genus responsible for most extraintestinal human infections.

Treatment of these organisms has become a major problem because of resistance to multiple antibiotics and potential transfer of plasmids to other organisms. Proper hand washing is crucial to prevent transmission from patient to patient via medical personnel. Contact isolation should be used for patients colonized or infected with highly antibiotic-resistant *Klebsiella* strains. *Klebsiella ozaenae* and *Klebsiella rhinoscleromatis* are infrequent isolates that are subspecies of *K. pneumoniae*; however, each is associated with a unique spectrum of disease. *K. ozaenae* is associated with atrophic rhinitis, a condition called ozena, and purulent infections of the nasal mucous membranes. *K. rhinoscleromatis* causes the granulomatous disease rhinoscleroma, an infection of the respiratory mucosa, oropharynx, nose, and paranasal sinuses.

Antibiotics may be indicated if symptoms are prolonged and in systemic infections. Refer to the antimicrobial susceptibilities for treatment.

Providencia spp

Providencia spp. belong to the *Enterobacteriaceae* family. This gram-negative bacilli is considered dysbiotic in the amount of 3 - 4+. *P. alcalifaciens* has been described as a possible diarrhea-causing pathogen in travelers and children in developing countries. In 1996, *P. alcalifaciens* caused a large food-borne outbreak involving 290 patients in Japan. *P. rettgeri* and *P. stuartii* also have been implicated as causes of travelers' diarrhea with symptoms including diarrhea, vomiting, fever and abdominal pain. Vomiting was a characteristic complaint among patients with diarrhea caused by *P. rettgeri*. The routes of exposure to *Providencia* have not been firmly established, but contaminated beef is a suspected source. Like *Proteus*, *Providencia* has the ability to alkalize urine and to promote the formation of crystals and calculi.

Antibiotics may be indicated if symptoms are prolonged. Refer to the bacterial sensitivities to identify the most appropriate antimicrobial therapy.

Imbalanced Flora

Imbalanced flora are those bacteria that reside in the host gastrointestinal tract and neither injure nor benefit the host. Certain dysbiotic bacteria may appear under the imbalanced category if found at low levels because they are not likely pathogenic at the levels detected. Imbalanced bacteria are commonly more abundant in association with insufficiency dysbiosis, and/or a fecal pH more towards the alkaline end of the reference range (5.8 - 7.0). Treatment with antimicrobial agents is unnecessary unless bacteria appear under the dysbiotic category.

**Order:** 999999-9999**Client #:** 999999**Doctor:** Sample Doctor, MD

Doctors Data Inc

123 Main St.

St. Charles, IL 60174 USA

Patient: Sample Patient**Id:** 999999**Age:** 38 **DOB:** 00/00/1986**Sex:** Female**Sample Collection****Date/Time****Date Collected**

05/14/2025

Date Received

05/23/2025

Date Reported

06/02/2025

Specimens Collected

3

Microbiology continued...

Cultured Yeast

Small amounts of yeast (+1) may be present in a healthy GI tract. However higher levels of yeast (> +1) are considered to be dysbiotic. A positive yeast culture and sensitivity to prescriptive and natural agents may help guide decisions regarding potential therapeutic intervention for yeast overgrowth. When investigating the presence of yeast, disparity may exist between culturing and microscopic examination. Yeast grows in colonies and is typically not uniformly dispersed throughout the stool. Further, some yeast may not survive transit through the intestines rendering it unviable for culturing. This may lead to undetectable or low levels of yeast identified by culture, despite a significant amount of yeast visualized microscopically. Therefore, both microscopic examination and culture are helpful in determining if abnormally high levels of yeast are present.

Dysbiotic Yeast

Yeast was cultured from this stool specimen at a level that is considered to be dysbiotic. A positive yeast culture and sensitivity to prescriptive and natural agents may help guide decisions regarding potential therapeutic intervention for chronic yeast syndrome. When investigating the presence of yeast, disparity may exist between culturing and microscopic examination. Yeast grows in colonies and is typically not uniformly dispersed throughout the stool. This may lead to undetectable or low levels of yeast identified by culture, despite a significant amount of yeast visualized microscopically.

GI Pathogens

Introduction

The GI Pathogen profile is performed using an FDA-cleared multiplex PCR system. It should be noted that PCR testing is much more sensitive than traditional techniques and allows for the detection of extremely low numbers of pathogens. PCR testing does not differentiate between viable and non-viable pathogens and should not be repeated until 21 days after completion of treatment or resolution to prevent false positives due to lingering traces of DNA. PCR testing can detect multiple pathogens in the patient's stool but does not differentiate the causative pathogen. All decisions regarding the need for treatment should take the patient's complete clinical history and presentation into account.

Parasitology

Microscopic yeast

Microscopic examination has revealed more yeast in this sample than normal. While small quantities of yeast (reported as rare) may be normal, yeast observed in higher amounts (moderate to many) is considered abnormal. Yeast does not appear to be dispersed uniformly throughout the stool. Yeast may therefore be observed microscopically, but not grow out on culture even when collected from the same bowel movement. Further, some yeast may not survive transit through the intestines rendering it unviable for culturing. Therefore, both microscopic examination and culture are helpful in determining if abnormally high levels of yeast are present. If significant yeast are reported by microscopy, but not by culture, consider the presentation of patient symptoms.

Stool Chemistries

Lactoferrin

The level of fecal lactoferrin is elevated in this sample. Lactoferrin is a biomarker of serious gastrointestinal inflammation which may be associated with inflammatory bowel disease (IBD) such as Ulcerative colitis (UC) or Crohn's disease (CD), but NOT Irritable bowel syndrome (IBS). Such distinction is critical because, although both IBD and IBS may share some common symptoms such as diarrhea, abdominal cramping and weight loss, the diseases are treated quite differently. IBD may become life threatening, requires lifelong treatment and possibly surgery. Very elevated lactoferrin should be reassessed in about four weeks, and if confirmed referral to a gastroenterologist should be considered. Lactoferrin is commonly high in breast-fed infants due to the high content in breast milk.

Patients with IBD oscillate between active and inactive disease states, and fecal lactoferrin levels increase 2-3 weeks prior to onset of clinical symptoms. During remission and effective treatment, fecal lactoferrin decreases significantly. Therefore, disease activity and efficacy of treatment can be monitored by following fecal lactoferrin levels. The test can be ordered separately to track disease activity in patients with IBD.

Moderately elevated levels of fecal lactoferrin can occur, often with fecal red and/or white blood cells, in association with invasive enteropathogens. Therefore, with moderately elevated levels of fecal lactoferrin, one should check for the presence of enteropathogens (eg. *Shigella*, *Campylobacter*, *Vibrio cholerae*, *Yersinia*).



Order: 999999-9999



Client #: 999999

Doctor: Sample Doctor, MD

Doctors Data Inc

123 Main St.

St. Charles, IL 60174 USA

Patient: Sample Patient

Id: 999999

Age: 38 **DOB:** 00/00/1986

Sex: Female

Sample Collection

Date/Time

Date Collected

05/14/2025

Date Received

05/23/2025

Date Reported

06/02/2025

Specimens Collected 3

Stool Chemistries continued...

Lysozyme

The level of lysozyme is elevated in this sample. Lysozyme is a biomarker of an inflammatory immune response in the gut. Moderate elevations in lysozyme are commonly associated with significant overgrowth of enteropathogens such as yeast, dysbiotic or pathogenic bacteria. Markedly elevated levels of lysozyme may occur with inflammatory bowel disease (IBD), such as Crohn's disease and Ulcerative colitis as well as other non-IBD intestinal diseases with diarrhea. If lysozyme is markedly elevated check the levels of calprotectin and lactoferrin. If either or both are very elevated reassess the levels in about four weeks. Lysozyme is commonly elevated for actively breast-feeding infants due to high maternal milk content.

Lysozyme is helpful in the determination of pathogen-induced inflammatory activity rather than IBD. Slightly-to moderately elevated levels of lysozyme may be remediated with elimination of an offending enteroinvasive microorganism and use of anti-inflammatory nutraceuticals.

Calprotectin (Moderately high)

The level of calprotectin is moderately elevated in this specimen. Moderate levels of calprotectin are an indicator of chronic inflammation. Inflammation at this level may be due to IBD in remission or inflammation caused by non-steroidal anti-inflammatories (NSAIDs).

Chronic inflammation of the gastrointestinal mucosa contributes to symptoms of IBD. Chronic stress is also known to contribute to symptom flare-ups and increased inflammation in IBD patients. Liver disease or the use of aspirin or nonsteroidal anti-inflammatory (NSAID) medications may variably elevate calprotectin levels.

Levels should be reassessed after about 4 weeks.

Short Chain Fatty Acids (SCFAs)

The total concentration and/or percentage distribution of the primary short chain fatty acids (SCFAs) are abnormal in this specimen. Beneficial bacteria that ferment non-digestible soluble fiber produce SCFAs that are pivotal in the regulation of intestinal health and function. Restoration of microbial abundance and diversity, and adequate daily consumption of soluble fiber and polyphenols can improve SCFA status.

The primary SCFAs butyrate, propionate and acetate are produced by predominant commensal bacteria via fermentation of soluble dietary fiber and intestinal mucus glycans. Key producers of SCFAs include *Faecalibacterium prausnitzii*, *Akkermansia muciniphila*, *Bacteroides fragilis*, *Bifidobacterium*, *Clostridium* and *Lactobacillus* spp. The SCFAs provide energy for intestinal cells, and regulate the actions of specialized mucosal cells that produce anti-inflammatory and antimicrobial factors, mucins that constitute the mucus barriers, and gut active peptides that facilitate appetite regulation and euglycemia. The SCFAs also contribute to a more acidic and anaerobic microenvironment that disfavors dysbiotic bacteria and yeast. Abnormal SCFAs may be associated with dysbiosis (including insufficiency dysbiosis), compromised intestinal barrier function (intestinal permeability) and inappropriate immune and inflammatory conditions.

"Seeding" with supplemental probiotics may contribute to improved production and status of SCFAs, but it is imperative to "feed" the beneficial microbes. Sources of soluble fiber that are available to the microbes include chick peas, beans, lentils, oat and rice bran, fructo- and galacto- oligosaccharides, and inulin.

β -glucuronidase

β -glucuronidase (β -G) is an enzyme that breaks the tight bond between glucuronic acid and toxins in the intestines. The liver and intestine bind toxins, steroid hormones and some dietary components to glucuronic acid. That is a protective process that limits absorption and enterohepatic resorption of toxins, and enhances excretion. A high level of activity of β -G in the gut is not desirable. A low level of β -G activity is not known to be of any direct clinical consequence.

β -glucuronidase is produced by the intestinal epithelium and many species of intestinal bacteria. Observational studies have indicated a correlation between high β -G activity and certain cancers, but a definitive causal relationship has not been established. Higher levels of β -G have been associated with higher circulating estrogens and lower fecal excretion of estrogens in premenopausal women. A potential dietary carcinogen derived from grilled/smoked meat and fish induces high β -G activity and prolongs internal exposure to the toxin in an experimental animal model.

**Order:** 999999-9999**Client #:** 999999**Doctor:** Sample Doctor, MD

Doctors Data Inc

123 Main St.

St. Charles, IL 60174 USA

Patient: Sample Patient**Id:** 999999**Age:** 38 **DOB:** 00/00/1986**Sex:** Female**Sample Collection****Date/Time****Date Collected**

05/14/2025

Date Received

05/23/2025

Date Reported

06/02/2025

Specimens Collected

3

Stool Chemistries continued...

Diet and intestinal bacterial imbalance modulate β -G activity. High fat, high protein and low fiber diets are associated with higher β -G activity compared to vegetarian or high soluble fiber diets. Higher β -G may be associated with an imbalanced intestinal microbiota profile. Some major bacterial producers of fecal β -G include *Bifidobacterium*, *Lactobacillus*, *Escherichia coli*, *Clostridium*, *Bacteroides fragilis* and other *Bacteroides* spp., *Mediterraneibacter gnavus*, and species that belong to the genera *Staphylococcus* and *Eubacterium*.

Low β -G activity is an indicator of abnormal metabolic activity among the intestinal microbiota that may be influenced by dietary extremes, diminished abundance and diversity of the intestinal microbiota, or heavy probiotic and/or prebiotic supplementation. A low fat, low meat and high fiber diet, such as consumed by strict vegetarians, may be benignly associated with lower β -G activity compared to a typical "Western diet." High-end consumption of soluble fiber (e.g. inulin) and supplementation with *Lactobacillus acidophilus* may be inconsequentially associated with lower fecal β -G.